

Plasma Dispersion Function for the Kappa Distribution

J. J. Podesta

The NASA STI Program Office ... in Profile

Since its founding, NASA has been dedicated to the advancement of aeronautics and space science. The NASA Scientific and Technical Information (STI) Program Office plays a key part in helping NASA maintain this important role.


The NASA STI Program Office is operated by Langley Research Center, the lead center for NASA's scientific and technical information. The NASA STI Program Office provides access to the NASA STI Database, the largest collection of aeronautical and space science STI in the world. The Program Office is also NASA's institutional mechanism for disseminating the results of its research and development activities. These results are published by NASA in the NASA STI Report Series, which includes the following report types:

- **TECHNICAL PUBLICATION.** Reports of completed research or a major significant phase of research that present the results of NASA programs and include extensive data or theoretical analysis. Includes compilations of significant scientific and technical data and information deemed to be of continuing reference value. NASA's counterpart of peer-reviewed formal professional papers but has less stringent limitations on manuscript length and extent of graphic presentations.
- **TECHNICAL MEMORANDUM.** Scientific and technical findings that are preliminary or of specialized interest, e.g., quick release reports, working papers, and bibliographies that contain minimal annotation. Does not contain extensive analysis.
- **CONTRACTOR REPORT.** Scientific and technical findings by NASA-sponsored contractors and grantees.
- **CONFERENCE PUBLICATION.** Collected papers from scientific and technical conferences, symposia, seminars, or other meetings sponsored or cosponsored by NASA.
- **SPECIAL PUBLICATION.** Scientific, technical, or historical information from NASA programs, projects, and mission, often concerned with subjects having substantial public interest.
- **TECHNICAL TRANSLATION.** English-language translations of foreign scientific and technical material pertinent to NASA's mission.

Specialized services that complement the STI Program Office's diverse offerings include creating custom thesauri, building customized databases, organizing and publishing research results . . . even providing videos.

For more information about the NASA STI Program Office, see the following:

- Access the NASA STI Program Home Page at <http://www.sti.nasa.gov/STI-homepage.html>
- E-mail your question via the Internet to help@sti.nasa.gov
- Fax your question to the NASA Access Help Desk at (301) 621-0134
- Telephone the NASA Access Help Desk at (301) 621-0390
- Write to:
NASA Access Help Desk
NASA Center for AeroSpace Information
7121 Standard Drive
Hanover, MD 21076-1320



Plasma Dispersion Function for the Kappa Distribution

John J. Podesta,

Courant Institute of Mathematical Sciences, New York University, NY, and

NASA Goddard Space Flight Center, Greenbelt, MD

National Aeronautics and
Space Administration
Goddard Space Flight Center
Greenbelt, Maryland 20771

Available from:

NASA Center for AeroSpace Information
7121 Standard Drive
Hanover, MD 21076-1320
Price Code: A17

National Technical Information Service
5285 Port Royal Road
Springfield, VA 22161
Price Code: A10

Abstract. The plasma dispersion function is computed for a homogeneous isotropic plasma in which the particle velocities are distributed according to a Kappa distribution. An ordinary differential equation is derived for the plasma dispersion function and it is shown that the solution can be written in terms of Gauss' hypergeometric function in the form $Z_\kappa(\xi) = i\kappa^{-3/2}(\kappa - 1/2)F(1, 2\kappa, \kappa + 1, t)$, where $t = (1 + i\xi/\sqrt{\kappa})/2$. Using the extensive theory of the hypergeometric function, various mathematical properties of the plasma dispersion function are derived including symmetry relations, series expansions, integral representations, and closed form expressions for integer and half-integer values of κ .

1 Introduction

Particle distribution functions with power law tails are frequently observed in space plasmas throughout the solar system. In practice, they are often modeled using Kappa distributions (defined below). The existence of such nonthermal, non-Maxwellian distribution functions can have profound effects on wave propagation and transport processes in these plasmas; effects that are of significant interest in space physics. The kinetic theory of such processes depends in an essential way on the plasma dielectric function and, therefore, on the plasma dispersion function.

The purpose of this paper is to derive the plasma dispersion function for the Kappa distribution in a new way and to investigate its mathematical properties. This function was derived independently by Summers and Thorne¹ and by Chateau and Meyer-Vernet² for the special case where $\kappa = 1, 2, 3, \dots$. It was later derived in the more general case $\kappa > 0$ by Mace and Hellberg.³ These authors all used the technique of contour integration. The approach taken here is different, being based instead on differential equations.

In order to be as consistent as possible with the formulation of the classical plasma dispersion function, the definition given here is slightly different from that of Summers and Thorne¹ and Mace and Hellberg.³ To clarify the reasons for doing this, the definition of the plasma dispersion function is briefly derived from first principles. The resulting dispersion relation for plasma waves, equation (29), is somewhat simplified in the present formulation.

Section 2 gives a concise list of the main results of this paper. The definition of the Kappa distribution is given in Section 3, and the definition of the plasma dispersion function is given in Section 4. In Section 5, the differential equation for the plasma dispersion function is derived and the solution is expressed in terms of Gauss' hypergeometric function. Symmetry relations and series expansions are derived in Sections 6 and 7. Integral representations are derived in Sections 8 and 9. Closed form expressions for integer and half-integer values of κ are derived in Sections 9 and 10, respectively.

2 Summary of Results

The important properties of the plasma dispersion function are listed here for easy reference. These properties are investigated in detail in the sections that follow. Throughout this paper the power function z^α , $\alpha \in \mathbb{R}$, is defined by its principal branch, that is, $z^\alpha = r^\alpha \exp(i\alpha\theta)$, where $z = r \exp(i\theta)$, $r \geq 0$, and $-\pi < \theta \leq \pi$.

Plasma Dispersion Function

$$Z_\kappa(\xi) = \frac{i(\kappa - \frac{1}{2})}{\kappa^{3/2}} F\left[1, 2\kappa, \kappa + 1, \frac{1}{2} \left(1 + \frac{i\xi}{\sqrt{\kappa}}\right)\right], \quad \kappa > 0. \quad (1)$$

If $\kappa = n$ is an integer, this function has a single pole of order n at the point $\xi = -i\sqrt{\kappa}$. Otherwise, it is analytic with a branch cut from $\xi = -i\sqrt{\kappa}$ to $-i\infty$ along the negative imaginary axis.

Behavior Near $\xi = -i\sqrt{\kappa}$

$$Z_\kappa(\xi) \sim \frac{i\pi^{1/2}}{2^{\kappa-1}\sqrt{\kappa}} \frac{\Gamma(\kappa)}{\Gamma(\kappa - \frac{1}{2})} \left(1 - \frac{i\xi}{\sqrt{\kappa}}\right)^{-\kappa} \quad \text{as } \xi \rightarrow -i\sqrt{\kappa}. \quad (2)$$

Symmetry Relations

$$Z_\kappa(-\xi^*) = -[Z_\kappa(\xi)]^*. \quad (3)$$

$$Z_\kappa(\xi) + Z_\kappa(-\xi) = \frac{2i\sqrt{\pi}}{\kappa^{1/2}} \frac{\Gamma(\kappa)}{\Gamma(\kappa - 1/2)} \left(1 + \frac{\xi^2}{\kappa}\right)^{-\kappa}. \quad (4)$$

Integral Representations

$$Z_\kappa(\xi) = \frac{1}{\sqrt{\pi\kappa}} \frac{\Gamma(\kappa)}{\Gamma(\kappa - \frac{1}{2})} \int_{-\infty}^{\infty} \frac{1}{(1+x^2)^\kappa} \cdot \frac{1}{(x-z)} dx, \quad \text{Im}(z) > 0, \quad (5)$$

where $z = \xi/\sqrt{\kappa}$,

$$Z_\kappa(\xi) = -\frac{2\kappa-1}{\kappa^{1/2}} (1+z^2)^{-\kappa} \int_i^z (1+t^2)^{\kappa-1} dt, \quad z \neq -i. \quad (6)$$

Differential Equation

$$(\xi^2 + \kappa)Z_\kappa'' + 2(\kappa+1)\xi Z_\kappa' + 2\kappa Z_\kappa = 0, \quad (7)$$

$$Z_\kappa(i\sqrt{\kappa}) = i \frac{\kappa - 1/2}{\kappa^{3/2}}, \quad Z_\kappa'(i\sqrt{\kappa}) = -\frac{\kappa - 1/2}{\kappa(\kappa+1)}. \quad (8)$$

Differentiation Formula

$$\frac{dZ_\kappa}{d\xi} = -\frac{2\kappa-1}{\kappa} \left[1 + \left(\frac{\kappa+1}{\kappa} \right)^{1/2} \xi Z_{\kappa+1}(\xi) \right]. \quad (9)$$

Closed Form Expression for $\kappa = n = 1, 2, 3 \dots$

$$Z_n(\xi) = i \frac{(n - \frac{1}{2})}{n^{1/2}} \frac{\Gamma(n)}{\Gamma(2n)} \sum_{m=0}^{n-1} \frac{\Gamma(m+n)}{m!} \left(\frac{1-iz}{2} \right)^{m-n}, \quad (10)$$

where $z = \xi/\sqrt{n}$.

Closed Form Expression for $\kappa = n + \frac{1}{2}$

$$Z_\kappa(\xi) = \frac{1}{(n + \frac{1}{2})^{1/2}} \frac{\Gamma(n + \frac{1}{2})}{\Gamma(n)} \left\{ \sum_{m=1}^n \frac{\Gamma(n+1-m)}{\Gamma(n + \frac{3}{2} - m)} z(1+z^2)^{-m} \right. \\ \left. - \frac{2}{\sqrt{\pi}} (1+z^2)^{-n-1/2} \left[\log(z + \sqrt{z^2+1}) - \frac{\pi i}{2} \right] \right\}, \quad (11)$$

where $z = \xi/\sqrt{\kappa}$ and $n = 1, 2, 3, \dots$

Series Expansion for $|z| < 1$

$$Z_\kappa(\xi) = i \frac{\sqrt{\pi}}{\kappa^{1/2}} \frac{\Gamma(\kappa)}{\Gamma(\kappa - \frac{1}{2})} (1 + z^2)^{-\kappa} - \frac{2\kappa - 1}{\kappa^{1/2}} z \left\{ 1 - \left(\frac{2\kappa + 1}{3} \right) z^2 + \left(\frac{2\kappa + 1}{3} \right) \left(\frac{2\kappa + 3}{5} \right) z^4 - \dots \right\}, \quad (12)$$

where $z = \xi/\sqrt{\kappa}$.

Series Expansion for $|z| > 1$

$$Z_\kappa(\xi) = \frac{i\sqrt{\pi}}{\sqrt{\kappa}} \frac{\Gamma(\kappa)}{\Gamma(\kappa - \frac{1}{2})} \left\{ 1 + i \tan(\pi\kappa) \operatorname{sgn}[\operatorname{Re}(z)] \right\} (z^2)^{-\kappa} \left(1 + \frac{1}{z^2} \right)^{-\kappa} - \frac{1}{\sqrt{\kappa}} \frac{1}{z} \left\{ 1 + \left(\frac{1}{2\kappa - 3} \right) \frac{1}{z^2} + \left(\frac{1}{2\kappa - 3} \right) \left(\frac{3}{2\kappa - 5} \right) \frac{1}{z^4} + \dots \right\}, \quad (13)$$

where $z = \xi/\sqrt{\kappa}$, $\kappa \neq \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots$, and the function $\operatorname{sgn}[\operatorname{Re}(z)]$ is defined by equation (81).

3 Kappa Distribution

In the astrophysics and space physics literature, the isotropic Kappa distribution in three dimensions is defined by

$$f_\kappa(\mathbf{v}) = A_\kappa \left(1 + \frac{v^2}{\kappa v_0^2} \right)^{-(\kappa+1)}, \quad \kappa > 1/2, \quad (14)$$

where $\mathbf{v} = (v_x, v_y, v_z)$ is the velocity vector, $v = |\mathbf{v}|$, and v_0 is some characteristic velocity. The normalization constant is

$$A_\kappa = \frac{1}{(\pi \kappa v_0^2)^{3/2}} \cdot \frac{\Gamma(\kappa + 1)}{\Gamma(\kappa - 1/2)} = \frac{1}{(\pi v_0^2)^{3/2} \sqrt{\kappa}} \cdot \frac{\Gamma(\kappa)}{\Gamma(\kappa - 1/2)} \quad (15)$$

and is chosen such that

$$4\pi \int_0^\infty f_\kappa(v) v^2 dv = 1. \quad (16)$$

The condition $\kappa > 1/2$ is necessary for this integral to converge. The Kappa distribution is closely related to the Beta distribution. In fact, the Beta function arises when computing the normalization factor A_κ and the statistical moments. In the limit as $\kappa \rightarrow \infty$, the Kappa distribution approaches the well known Maxwell distribution e^{-v^2/v_0^2} .

The velocity moments of the Kappa distribution are given by

$$\langle v^n \rangle = 4\pi \int_0^\infty f_\kappa(v) v^{n+2} dv = \frac{2}{\sqrt{\pi}} (\kappa v_0^2)^{n/2} \frac{\Gamma(\frac{n+3}{2}) \Gamma(\kappa - \frac{n+1}{2})}{\Gamma(\kappa - \frac{1}{2})}, \quad (17)$$

where n is an integer and $0 \leq n \leq 2(\kappa - 1)$. For an arbitrary real power $n \geq 0$, this integral is finite if and only if $n < 2\kappa - 1$. In applications, the equivalent temperature of an isotropic Kappa distribution is defined by the relation

$$\frac{1}{2} m \langle v^2 \rangle = \frac{3}{2} k_B T \quad (18)$$

which implies

$$k_B T = \frac{1}{2} m v_0^2 \left(\frac{\kappa}{\kappa - 3/2} \right), \quad (19)$$

where k_B is Boltzmann's constant. The second moment $\langle v^2 \rangle$ is finite if and only if $\kappa > 3/2$.

4 Plasma Dispersion Function

In the Vlasov theory of plasma waves⁴ the propagation of electrostatic waves (having $\mathbf{E} \parallel \mathbf{k}$) in a homogeneous isotropic plasma is governed by the dispersion relation $\varepsilon_L(\mathbf{k}, s) = 0$, where

$$\varepsilon_L(\mathbf{k}, s) = 1 - \sum_\alpha \frac{\omega_{p\alpha}^2}{k^2} \int_{-\infty}^{\infty} \frac{\partial F_{\alpha 0} / \partial u}{u - is/|k|} du \quad (20)$$

is the longitudinal dielectric function and SI units are used throughout. The index α denotes the different particle species, $\omega_{p\alpha}$ is the plasma frequency for species α , \mathbf{k} is the Fourier wavevector with magnitude $k = |\mathbf{k}|$, s is the Laplace transform variable, and $F_{\alpha 0}$ is related to the particle distribution function $f_{\alpha 0}$ by

$$F_{\alpha 0}(u) = \int f_{\alpha 0}(\mathbf{v}) \delta\left(u - \frac{\mathbf{k} \cdot \mathbf{v}}{|k|}\right) d\mathbf{v}. \quad (21)$$

The distribution function $f_{\alpha 0}(\mathbf{v})$ is normalized to unity.

Spatial isotropy implies $f_0(\mathbf{v}) = f_0(v^2)$, where $v = |\mathbf{v}|$. After a rotation of coordinates which aligns the v_z direction with \mathbf{k} , the last equation becomes

$$F_{\alpha 0}(u) = \int \int f_{\alpha 0}(v_x^2 + v_y^2 + u^2) dv_x dv_y \quad (22)$$

$$= 2\pi \int_{|u|}^{\infty} f_{\alpha 0}(v^2) v dv \quad (23)$$

and, therefore,

$$\frac{\partial F_{\alpha 0}}{\partial u} = -2\pi u f_{\alpha 0}(u^2). \quad (24)$$

It is usually the case that $f_{\alpha 0}$ has the functional form $f_{\alpha 0}(v/v_{\alpha 0})$, where $v_{\alpha 0}$ is some characteristic velocity; consequently, $F_{\alpha 0} = F_{\alpha 0}(u/v_{\alpha 0})$. Changing to normalized velocity variables:

$$u' = \frac{u}{v_{\alpha 0}}, \quad \tilde{F}(u') = v_{\alpha 0} F(u'), \quad \frac{\partial F}{\partial u} = \frac{1}{v_{\alpha 0}^2} \frac{\partial \tilde{F}}{\partial u'}, \quad (25)$$

the dispersion relation (20) can be written

$$1 - \sum_{\alpha} \frac{\omega_{p\alpha}^2}{k^2 v_{\alpha 0}^2} Z' \left(\alpha, \frac{is}{|k|v_{\alpha 0}} \right) = 0, \quad (26)$$

where, after dropping the primes on u ,

$$Z'(\alpha, \xi) = \int_{-\infty}^{\infty} \frac{\partial \tilde{F}_{\alpha 0}/\partial u}{u - \xi} du = \int_{-\infty}^{\infty} \frac{\tilde{F}_{\alpha 0}(u)}{(u - \xi)^2} du = \frac{d}{d\xi} \int_{-\infty}^{\infty} \frac{\tilde{F}_{\alpha 0}(u)}{u - \xi} du \quad (27)$$

and, by definition,

$$Z(\alpha, \xi) = \int_{-\infty}^{\infty} \frac{\tilde{F}_{\alpha 0}(u)}{u - \xi} du, \quad \text{Im}(\xi) > 0, \quad (28)$$

is the *plasma dispersion function*. It is defined for $\text{Im}(\xi) \leq 0$ by analytic continuation. Introducing the complex frequency $\omega = is$ which is consistent with Fourier modes of the form $e^{i(\mathbf{k} \cdot \mathbf{x} - \omega t)}$, the plasma dispersion relation takes the standard form⁵

$$1 - \sum_{\alpha} \frac{\omega_{p\alpha}^2}{k^2 v_{\alpha 0}^2} Z' \left(\frac{\omega}{|k|v_{\alpha 0}} \right) = 0. \quad (29)$$

If $f_{\alpha 0}$ is a Maxwell distribution

$$f_{\alpha 0}(v) = \frac{1}{(\pi v_0^2)^{3/2}} e^{-v^2/v_0^2}, \quad (30)$$

then the evaluation of (23) and (28) yields the well known plasma dispersion function⁶

$$Z(\xi) = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \frac{e^{-u^2}}{u - \xi} du, \quad \text{Im}(\xi) > 0. \quad (31)$$

If $f_{\alpha 0}$ is a Kappa distribution (14), then the evaluation of (23) and (28) yields

$$F_{\alpha 0}(u) = \frac{1}{\sqrt{\pi \kappa} v_0} \cdot \frac{\Gamma(\kappa)}{\Gamma(\kappa - \frac{1}{2})} \cdot \frac{1}{(1 + u^2/\kappa v_0^2)^\kappa} \quad (32)$$

and

$$Z_\kappa(\xi) = \frac{1}{\sqrt{\pi \kappa}} \cdot \frac{\Gamma(\kappa)}{\Gamma(\kappa - \frac{1}{2})} \int_{-\infty}^{\infty} \frac{1}{(u - \xi)(1 + u^2/\kappa)^\kappa} du. \quad (33)$$

This is the dispersion function for the isotropic Kappa distribution. It generalizes the well known result (31) for the isotropic Maxwell distribution. It is different from the definition of Summers and Thorne¹ which was also adopted by Mace and Hellberg.³ These authors employ an exponent $\kappa + 1$ rather than κ in the integrand. Consequently, the function $Z_\kappa(\xi)$ of Summers and Thorne is more closely related to the derivative $Z'_\kappa(\xi)$ of the function defined here.

5 Representation by Gauss' Hypergeometric Function

For purposes of mathematical manipulation, it is convenient to let $x = u/\sqrt{\kappa}$ in equation (33) and write

$$Z_\kappa(\xi) = \frac{1}{\sqrt{\pi \kappa}} \frac{\Gamma(\kappa)}{\Gamma(\kappa - \frac{1}{2})} \int_{-\infty}^{\infty} \frac{1}{(1 + x^2)^\kappa} \cdot \frac{1}{(x - z)} dx, \quad \text{Im}(z) > 0, \quad (34)$$

where $z = \xi/\sqrt{\kappa}$. Therefore, the calculation of the plasma dispersion function reduces to the evaluation of the integral

$$I = \int_{-\infty}^{\infty} \frac{1}{(1 + x^2)^\kappa} \cdot \frac{1}{(x - z)} dx, \quad \text{Im}(z) > 0. \quad (35)$$

It is easy to see that this integral converges for all $\kappa > 0$. There are two cases to consider: the case where $\kappa > 0$ is an integer, and the case when $\kappa > 0$ is not an integer. The solution in the first case was derived independently by Chateau and Meyer-Vernet² and by Summers and Thorne.¹ The solution in the second case was derived by Mace and Hellberg.³ As will now be shown, the solution for arbitrary $\kappa > 0$ is easily obtained by first finding the differential equation for $I(z)$. This method of approach is an alternative to the method of contour integration used by the above authors.

Differentiating equation (35) with respect to z yields

$$I'(z) = \int_{-\infty}^{\infty} \frac{1}{(1+x^2)^\kappa} \cdot \frac{1}{(x-z)^2} dx. \quad (36)$$

Multiplying this by z and then using the identity

$$1 + \frac{z}{x-z} = \frac{x}{x-z}, \quad (37)$$

it follows that

$$zI' + I = \int_{-\infty}^{\infty} \frac{1}{(1+x^2)^\kappa} \cdot \frac{x}{(x-z)^2} dx. \quad (38)$$

Differentiating this equation with respect to z , multiplying the result by z , and then using the identity (37) yields

$$z^2 I'' + 4zI' + 2I = \int_{-\infty}^{\infty} \frac{1}{(1+x^2)^\kappa} \cdot \frac{2x^2}{(x-z)^3} dx. \quad (39)$$

Now differentiate (36) and add the result to the last equation to obtain

$$(1+z^2)I'' + 4zI' + 2I = \int_{-\infty}^{\infty} \frac{1}{(1+x^2)^{\kappa-1}} \cdot \frac{2}{(x-z)^3} dx. \quad (40)$$

Integrating the remaining integral by parts and comparing the result with equation (38), the right hand side of (40) equals

$$-2(\kappa-1) \int_{-\infty}^{\infty} \frac{1}{(1+x^2)^\kappa} \cdot \frac{x}{(x-z)^2} dx = -2(\kappa-1)(zI' + I). \quad (41)$$

Thus, the differential equation for $I(z)$ is

$$(1+z^2)I'' + 2(\kappa+1)zI' + 2\kappa I = 0. \quad (42)$$

To obtain a unique solution, it is necessary to impose the initial conditions

$$I(z = i) = I_0, \quad (43)$$

$$I'(z = i) = \frac{2\kappa}{\kappa + 1} \frac{iI_0}{2}, \quad (44)$$

where I_0 is evaluated in Appendix A with the result

$$I_0 = i\sqrt{\pi} \frac{\Gamma(\kappa + \frac{1}{2})}{\Gamma(\kappa + 1)}. \quad (45)$$

Making the transformation

$$t = \frac{1 + iz}{2} \quad (46)$$

the differential equation (42) takes the form

$$t(1 - t)I'' + (\kappa + 1)(1 - 2t)I' - 2\kappa I = 0, \quad (47)$$

where

$$I(t = 0) = I_0, \quad (48)$$

$$I'(t = 0) = \frac{2\kappa}{\kappa + 1} I_0. \quad (49)$$

Equation (47) is now in the standard form of the well known hypergeometric equation.

Recall that the solution of the hypergeometric equation

$$z(1 - z)F'' + [c - (a + b + 1)z]F' - abF = 0 \quad (50)$$

with initial conditions

$$F(0) = 1, \quad (51)$$

$$F'(0) = \frac{ab}{c} \quad (52)$$

is given by Gauss' hypergeometric function $F(a, b, c, z) = {}_2F_1(a, b, c, z)$. This is a single valued analytic function of z in the complex plane with a branch cut from 1 to ∞ along the positive real axis. For fixed z it is analytic in a ,

b , and c separately with the exception of simple poles at $c = 0, -1, -2, \dots$. In the unit disk it has the series representation

$$F(a, b, c, z) = 1 + \frac{ab}{c} \frac{z}{1!} + \frac{a(a+1)b(b+1)}{c(c+1)} \frac{z^2}{2!} + \dots, \quad |z| < 1, \quad (53)$$

which also shows that it is symmetric in a and b . The series terminates if a or $b \in \{0, -1, -2, \dots\}$, in which case $F(a, b, c, z)$ is entire because it reduces to a polynomial in z . The properties of the function F have been thoroughly studied and documented. Therefore, for all intents and purposes it can be considered to be a closed form function in the same way that $\sin(z)$ and $\cos(z)$ are.

Choosing

$$a = 1, \quad b = 2\kappa, \quad c = \kappa + 1, \quad (54)$$

equation (47) is equivalent to the hypergeometric equation (50); hence, the unique solution of (47) is given by

$$I(t) = i\sqrt{\pi} \frac{\Gamma(\kappa + \frac{1}{2})}{\Gamma(\kappa + 1)} F(1, 2\kappa, \kappa + 1, t). \quad (55)$$

Using (46), a closed form expression for the integral (35) is

$$I(z) = i\sqrt{\pi} \frac{\Gamma(\kappa + \frac{1}{2})}{\Gamma(\kappa + 1)} F\left(1, 2\kappa, \kappa + 1, \frac{1 + iz}{2}\right), \quad (56)$$

which is valid for all $\kappa > 0$, in fact, for $\text{Re}(\kappa) > 0$. Substituting this into (34), the plasma dispersion function is found to be

$$Z_\kappa(\xi) = \frac{i(\kappa - \frac{1}{2})}{\kappa^{3/2}} F(1, 2\kappa, \kappa + 1, t), \quad (57)$$

where

$$t = \frac{1 + iz}{2} = \frac{1}{2} \left(1 + \frac{i\xi}{\sqrt{\kappa}}\right). \quad (58)$$

It is important to note that the function $Z_\kappa(\xi)$ inherits all the properties of $F(a, b, c, z)$. Because $F(a, b, c, z)$ has a branch cut from 1 to ∞ , the function $Z_\kappa(\xi)$ is single valued analytic with a branch cut from $\xi = -i\sqrt{\kappa}$ to $\xi = -i\infty$ along the negative imaginary axis. Here, the ξ -plane is the complex frequency

plane, $\xi = is/|k|v_0$, and the lower half plane consists of all the damped modes.

The behavior in the neighborhood of the singularity $\xi = -i\sqrt{\kappa}$ can be obtained from the identity⁷

$$F(a, b, c, z) = (1 - z)^{c-a-b} F(c - a, c - b, c, z). \quad (59)$$

This yields

$$F\left(1, 2\kappa, \kappa + 1, \frac{1 + iz}{2}\right) = \left(\frac{1 - iz}{2}\right)^{-\kappa} F\left(\kappa, 1 - \kappa, \kappa + 1, \frac{1 + iz}{2}\right). \quad (60)$$

Using the formula⁷

$$F(a, b, c, 1) = \frac{\Gamma(c)\Gamma(c - a - b)}{\Gamma(c - a)\Gamma(c - b)}, \quad c > a + b, \quad (61)$$

it follows that

$$\lim_{z \rightarrow -i} \left(\frac{1 - iz}{2}\right)^{\kappa} F\left(1, 2\kappa, \kappa + 1, \frac{1 + iz}{2}\right) = \frac{\Gamma(\kappa + 1)\Gamma(\kappa)}{\Gamma(1)\Gamma(2\kappa)} = \frac{\pi^{1/2}\Gamma(\kappa + 1)}{2^{2\kappa-1}\Gamma(\kappa + \frac{1}{2})}. \quad (62)$$

Therefore, one obtains the asymptotic relation

$$Z_{\kappa}(\xi) \sim \frac{i\pi^{1/2}}{2^{\kappa-1}\sqrt{\kappa}} \frac{\Gamma(\kappa)}{\Gamma(\kappa - \frac{1}{2})} \left(1 - \frac{i\xi}{\sqrt{\kappa}}\right)^{-\kappa} \quad \text{as } \xi \rightarrow -i\sqrt{\kappa}. \quad (63)$$

6 Symmetry Relations and Taylor Expansion

Using the well established theory of the hypergeometric function, the properties of $Z_{\kappa}(\xi)$ may be derived with little effort. From the series (53), it is obvious that for a , b , and c real,

$$F(a, b, c, z^*) = [F(a, b, c, z)]^*. \quad (64)$$

Therefore, replacing ξ with $-\xi^*$ in equations (57) and (58), it follows that

$$Z_{\kappa}(-\xi^*) = -[Z_{\kappa}(\xi)]^*. \quad (65)$$

This relates the values of $Z_\kappa(\xi)$ in the left half-plane to those in the right half-plane. The classical dispersion function for the Maxwell distribution has the same property.

From the identity^{8,9}

$$F\left(2a, 2b, a + b + \frac{1}{2}, \frac{1+z}{2}\right) = \frac{\Gamma(a + b + \frac{1}{2})\Gamma(\frac{1}{2})}{\Gamma(a + \frac{1}{2})\Gamma(b + \frac{1}{2})}F\left(a, b, \frac{1}{2}, z^2\right) \\ - z \frac{\Gamma(a + b + \frac{1}{2})\Gamma(-\frac{1}{2})}{\Gamma(a)\Gamma(b)}F\left(a + \frac{1}{2}, b + \frac{1}{2}, \frac{3}{2}, z^2\right) \quad (66)$$

one has

$$F\left(1, 2\kappa, \kappa + 1, \frac{1+iz}{2}\right) = \sqrt{\pi} \frac{\Gamma(\kappa + 1)}{\Gamma(\kappa + \frac{1}{2})}F\left(\frac{1}{2}, \kappa, \frac{1}{2}, -z^2\right) \\ + 2\kappa iz F\left(1, \kappa + \frac{1}{2}, \frac{3}{2}, -z^2\right). \quad (67)$$

Substituting the closed form expression⁷

$$F(a, b, b, z) = F(b, a, b, z) = (1 - z)^{-a}, \quad (68)$$

equation (67) takes the form

$$F\left(1, 2\kappa, \kappa + 1, \frac{1+iz}{2}\right) = \sqrt{\pi} \frac{\Gamma(\kappa + 1)}{\Gamma(\kappa + \frac{1}{2})}(1 + z^2)^{-\kappa} + 2\kappa iz F\left(1, \kappa + \frac{1}{2}, \frac{3}{2}, -z^2\right). \quad (69)$$

Multiplying this equation by the leading coefficient in equation (57) yields

$$Z_\kappa(\xi) = \frac{i\sqrt{\pi}}{\kappa^{1/2}} \frac{\Gamma(\kappa)}{\Gamma(\kappa - \frac{1}{2})}(1 + z^2)^{-\kappa} - \frac{2\kappa - 1}{\kappa^{1/2}} z F\left(1, \kappa + \frac{1}{2}, \frac{3}{2}, -z^2\right), \quad (70)$$

where $z = \xi/\sqrt{\kappa}$. Changing ξ to $-\xi$ (or z to $-z$) and adding the result to the last equation yields the symmetry relation

$$Z_\kappa(\xi) + Z_\kappa(-\xi) = \frac{2i\sqrt{\pi}}{\kappa^{1/2}} \frac{\Gamma(\kappa)}{\Gamma(\kappa - \frac{1}{2})} \left(1 + \frac{\xi^2}{\kappa}\right)^{-\kappa}. \quad (71)$$

This relates the values of $Z_\kappa(\xi)$ in the upper half-plane to those in the lower half-plane. In the limit as $\kappa \rightarrow \infty$ one recovers the symmetry relation

$$Z(\xi) + Z(-\xi) = 2i\sqrt{\pi}e^{-\xi^2} \quad (72)$$

for the Maxwellian dispersion function.

Using the series expansion (53) for $F(a, b, c, z)$, equation (70) immediately yields the following series expansion valid for $|z| < 1$:

$$Z_\kappa(\xi) = i \frac{\sqrt{\pi}}{\kappa^{1/2}} \frac{\Gamma(\kappa)}{\Gamma(\kappa - \frac{1}{2})} (1 + z^2)^{-\kappa} - \frac{2\kappa - 1}{\kappa^{1/2}} z \left\{ 1 + \left(\frac{2\kappa + 1}{3} \right) (-z^2) + \left(\frac{2\kappa + 1}{3} \right) \left(\frac{2\kappa + 3}{5} \right) (-z^2)^2 + \dots \right\}. \quad (73)$$

Here, the even powers are neatly summed into the first term. If necessary, the first term can be expanded using the binomial theorem. The series expansion can also be written in the more compact form

$$Z_\kappa(\xi) = i \frac{\sqrt{\pi}}{\kappa^{1/2}} \frac{\Gamma(\kappa)}{\Gamma(\kappa - \frac{1}{2})} (1 + z^2)^{-\kappa} - \frac{\sqrt{\pi}}{\kappa^{1/2}} \frac{z}{\Gamma(\kappa - \frac{1}{2})} \sum_{n=0}^{\infty} \frac{\Gamma(\kappa + n + 1/2)}{\Gamma(n + 3/2)} (-z^2)^n, \quad |z| < 1, \quad (74)$$

however, the coefficients in (73) are much more explicit.

7 Series Expansion for Large z

Begin with the identity⁷

$$F(a, b, c, z) = \frac{\Gamma(c)\Gamma(b-a)}{\Gamma(b)\Gamma(c-a)} (-z)^{-a} F\left(a, a-c+1, a-b+1, \frac{1}{z}\right) + \frac{\Gamma(c)\Gamma(a-b)}{\Gamma(a)\Gamma(c-b)} (-z)^{-b} F\left(b, b-c+1, b-a+1, \frac{1}{z}\right). \quad (75)$$

Applying this to the last term in equation (69) yields

$$F\left(1, 2\kappa, \kappa + 1, \frac{1 + iz}{2}\right) = \sqrt{\pi} \frac{\Gamma(\kappa + 1)}{\Gamma(\kappa + \frac{1}{2})} (1 + z^2)^{-\kappa} + \frac{i\kappa}{(\kappa - \frac{1}{2})} z^{-1} F\left(1, \frac{1}{2}, \frac{3}{2} - \kappa, -\frac{1}{z^2}\right) + i\sqrt{\pi}\kappa \frac{\Gamma(\frac{1}{2} - \kappa)}{\Gamma(1 - \kappa)} z(z^2)^{-(\kappa+1/2)} F\left(\kappa + \frac{1}{2}, \kappa, \kappa + \frac{1}{2}, -\frac{1}{z^2}\right). \quad (76)$$

In addition, using the identity (68), this simplifies to

$$\begin{aligned} F\left(1, 2\kappa, \kappa + 1, \frac{1 + iz}{2}\right) &= \sqrt{\pi} \frac{\Gamma(\kappa + 1)}{\Gamma(\kappa + \frac{1}{2})} (1 + z^2)^{-\kappa} \\ &\quad + \frac{i\kappa}{(\kappa - \frac{1}{2})} z^{-1} F\left(1, \frac{1}{2}, \frac{3}{2} - \kappa, -\frac{1}{z^2}\right) \\ &\quad + i\sqrt{\pi}\kappa \frac{\Gamma(\frac{1}{2} - \kappa)}{\Gamma(1 - \kappa)} z(z^2)^{-(\kappa+1/2)} \left(1 + \frac{1}{z^2}\right)^{-\kappa}. \end{aligned} \quad (77)$$

Using the reflection formula

$$\Gamma(z)\Gamma(1 - z) = \frac{\pi}{\sin(\pi z)}, \quad (78)$$

the last equation takes the form

$$\begin{aligned} F\left(1, 2\kappa, \kappa + 1, \frac{1 + iz}{2}\right) &= \sqrt{\pi} \frac{\Gamma(\kappa + 1)}{\Gamma(\kappa + \frac{1}{2})} (1 + z^2)^{-\kappa} \\ &\quad + \frac{i\kappa}{(\kappa - \frac{1}{2})} z^{-1} F\left(1, \frac{1}{2}, \frac{3}{2} - \kappa, -\frac{1}{z^2}\right) \\ &\quad + \sqrt{\pi} i \tan(\pi\kappa) \frac{\Gamma(\kappa + 1)}{\Gamma(\kappa + \frac{1}{2})} z(z^2)^{-(\kappa+1/2)} \left(1 + \frac{1}{z^2}\right)^{-\kappa}. \end{aligned} \quad (79)$$

In order to get the phases right special care must be taken when combining the different power functions. This is because, for the principal branch of the power function, it is not always true that $(z^\alpha)^\beta = z^{\alpha\beta}$ or $z_1^\alpha z_2^\alpha = (z_1 z_2)^\alpha$. However, using the identities

$$z^{\alpha+\beta} = z^\alpha z^\beta, \quad (1 + z)^\alpha = z^\alpha \left(1 + \frac{1}{z}\right)^\alpha, \quad (80)$$

and

$$z(z^2)^{-1/2} = \text{sgn}[\text{Re}(z)] \equiv \begin{cases} +1 & \text{if } \text{Re}(z) \geq 0 \text{ and } \arg(z) \neq -\pi/2 \\ -1 & \text{if } \text{Re}(z) \leq 0 \text{ and } \arg(z) \neq +\pi/2, \end{cases} \quad (81)$$

which are valid for all $z \neq 0$, equation (79) can be written

$$\begin{aligned} F\left(1, 2\kappa, \kappa + 1, \frac{1 + iz}{2}\right) &= \frac{i\kappa}{(\kappa - \frac{1}{2})} z^{-1} F\left(1, \frac{1}{2}, \frac{3}{2} - \kappa, -\frac{1}{z^2}\right) \\ &\quad + \sqrt{\pi} \frac{\Gamma(\kappa + 1)}{\Gamma(\kappa + \frac{1}{2})} \left\{ 1 + i \tan(\pi\kappa) \text{sgn}[\text{Re}(z)] \right\} (z^2)^{-\kappa} \left(1 + \frac{1}{z^2}\right)^{-\kappa}. \end{aligned} \quad (82)$$

It should be noted that for half-integer values of κ the right-hand side is undefined. Otherwise, this equation holds for all $z \neq 0$ in the complex plane with a branch cut from $-i$ to $-i\infty$ along the negative imaginary axis. For $|z| > 1$, the hypergeometric function on the right-hand side of (82) can be expanded using the series (53) to obtain the expansion

$$\begin{aligned} F\left(1, 2\kappa, \kappa + 1, \frac{1 + iz}{2}\right) = \\ \sqrt{\pi} \frac{\Gamma(\kappa + 1)}{\Gamma(\kappa + \frac{1}{2})} \left\{ 1 + i \tan(\pi\kappa) \operatorname{sgn}[\operatorname{Re}(z)] \right\} (z^2)^{-\kappa} \left(1 + \frac{1}{z^2} \right)^{-\kappa} \\ + \frac{i\kappa}{(\kappa - \frac{1}{2})} \frac{1}{z} \left\{ 1 + \frac{\frac{1}{2}}{\frac{3}{2} - \kappa} \left(-\frac{1}{z^2} \right) + \frac{\frac{1}{2} \cdot \frac{3}{2}}{(\frac{3}{2} - \kappa)(\frac{5}{2} - \kappa)} \left(-\frac{1}{z^2} \right)^2 + \dots \right\}, \quad (83) \end{aligned}$$

$|z| > 1$. The factor $\operatorname{sgn}[\operatorname{Re}(z)]$ ensures that the right-hand side is continuous throughout the upper half plane. This expansion can be arrived at independently using the theory of Laplace transforms as shown in Appendix B. Substituting (83) into (57) yields

$$\begin{aligned} Z_\kappa(\xi) = \frac{i\sqrt{\pi}}{\sqrt{\kappa}} \frac{\Gamma(\kappa)}{\Gamma(\kappa - \frac{1}{2})} \left\{ 1 + i \tan(\pi\kappa) \operatorname{sgn}[\operatorname{Re}(z)] \right\} (z^2)^{-\kappa} \left(1 + \frac{1}{z^2} \right)^{-\kappa} \\ - \frac{1}{\sqrt{\kappa}} \frac{1}{z} \left\{ 1 + \frac{\frac{1}{2}}{\frac{3}{2} - \kappa} \left(-\frac{1}{z^2} \right) + \frac{\frac{1}{2} \cdot \frac{3}{2}}{(\frac{3}{2} - \kappa)(\frac{5}{2} - \kappa)} \left(-\frac{1}{z^2} \right)^2 + \dots \right\}, \quad (84) \end{aligned}$$

which is valid for $|z| > 1$ and $\kappa \neq \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots$. It is important to note that this expansion is exact, not just asymptotic.

8 Integration of the Differential Equation

The differential equation (42) can be written in the symmetric form

$$\frac{d}{dz} \left\{ (1 + z^2)^{1-\kappa} \frac{d}{dz} \left[(1 + z^2)^\kappa I(z) \right] \right\} = 0. \quad (85)$$

This has the first integral

$$(1 + z^2)^{1-\kappa} \frac{d}{dz} \left[(1 + z^2)^\kappa I(z) \right] = C_\kappa, \quad (86)$$

where

$$C_\kappa = -2\kappa \int_{-\infty}^{\infty} \frac{1}{(1+x^2)^{\kappa+1}} dx = -2\sqrt{\pi} \frac{\Gamma(\kappa + \frac{1}{2})}{\Gamma(\kappa)}. \quad (87)$$

Equations (86) and (87) can be verified by substituting (35) into (86), carrying out the differentiation, and then using integration by parts and the identity (37). Therefore, the solution can be written

$$I(z) = C_\kappa (1+z^2)^{-\kappa} \int_i^z (1+t^2)^{\kappa-1} dt, \quad (88)$$

where the lower limit of integration is chosen such that

$$\lim_{z \rightarrow i} I(z) = i\sqrt{\pi} \frac{\Gamma(\kappa + \frac{1}{2})}{\Gamma(\kappa + 1)}. \quad (89)$$

If $\kappa = 1/2$, the integral (88) has the closed form solution

$$I(z) = -2(1+z^2)^{-1/2} \left[\sinh^{-1}(z) - \frac{\pi i}{2} \right], \quad (90)$$

where

$$\sinh^{-1}(z) = \int_0^z \frac{1}{(1+t^2)^{1/2}} dt. \quad (91)$$

Equivalently, this can be written

$$\sinh^{-1}(z) = \log \left(z + \sqrt{z^2 + 1} \right), \quad (92)$$

where $\log(z)$ is the principal branch of the logarithm. Substituting the result (88) into (34), the plasma dispersion function can be written

$$Z_\kappa(\xi) = -\frac{2\kappa-1}{\kappa^{1/2}} (1+z^2)^{-\kappa} \int_i^z (1+t^2)^{\kappa-1} dt, \quad (93)$$

where $z = \xi/\sqrt{\kappa}$. This holds in both the upper and lower half planes and therefore represents the analytic continuation of Z_κ .

9 Solutions for Half-Integer Values: $\kappa = n + \frac{1}{2}$

Setting $\kappa = n + \frac{1}{2}$ in equation (69) yields

$$F\left(1, 2\kappa, \kappa + 1, \frac{1+iz}{2}\right) = \sqrt{\pi} \frac{\Gamma(n + \frac{3}{2})}{\Gamma(n+1)} (1+z^2)^{-n-1/2} \\ + (2n+1)iz F\left(1, n+1, \frac{3}{2}, -z^2\right). \quad (94)$$

By repeated application of the identity⁷

$$(b-a)(1-z)F(a, b, c, z) = (c-a)F(a-1, b, c, z) - (c-b)F(a, b-1, c, z) \quad (95)$$

one may prove the formula

$$F\left(1, n+1, \frac{3}{2}, -z^2\right) = \frac{1}{2} \sum_{m=1}^n \frac{\Gamma(n+1-m)}{\Gamma(n+1)} \frac{\Gamma(n+\frac{1}{2})}{\Gamma(n+\frac{3}{2}-m)} (1+z^2)^{-m} \\ + \frac{1}{2} \frac{\Gamma(1)}{\Gamma(n+1)} \frac{\Gamma(n+\frac{1}{2})}{\Gamma(\frac{3}{2})} (1+z^2)^{-n} F\left(1, 1, \frac{3}{2}, -z^2\right), \quad (96)$$

where $n = 1, 2, 3, \dots$. Inserting this formula into (94) and using the closed form expression⁷

$$F\left(1, 1, \frac{3}{2}, -z^2\right) = \frac{1}{iz} (1+z^2)^{-1/2} \sin^{-1}(iz) \quad (97)$$

one finds

$$F\left(1, 2\kappa, \kappa + 1, \frac{1+iz}{2}\right) = \sqrt{\pi} \frac{\Gamma(n + \frac{3}{2})}{\Gamma(n+1)} (1+z^2)^{-n-1/2} \left[1 + \frac{2}{\pi} \sin^{-1}(iz) \right] \\ + \frac{\Gamma(n + \frac{3}{2})}{\Gamma(n+1)} \sum_{m=1}^n \frac{\Gamma(n+1-m)}{\Gamma(n+\frac{3}{2}-m)} iz (1+z^2)^{-m}, \quad (98)$$

where $\kappa = n + \frac{1}{2}$. Thus, from (56) one obtains the closed form solutions

$$I(z) = -2(1+z^2)^{-n-1/2} \left[\sinh^{-1}(z) - \frac{\pi i}{2} \right] \\ - \sqrt{\pi} \sum_{m=1}^n \frac{\Gamma(n+1-m)}{\Gamma(n+\frac{3}{2}-m)} z (1+z^2)^{-m}, \quad (99)$$

$n = 1, 2, 3, \dots$ The first few solutions are as follows; for $\kappa = \frac{3}{2}$:

$$I(z) = -2(1 + z^2)^{-3/2} \left[\sinh^{-1}(z) - \frac{\pi i}{2} \right] - 2z(1 + z^2)^{-1}, \quad (100)$$

for $\kappa = \frac{5}{2}$:

$$I(z) = -2(1 + z^2)^{-5/2} \left[\sinh^{-1}(z) - \frac{\pi i}{2} \right] - \frac{4}{3}z(1 + z^2)^{-1} - 2z(1 + z^2)^{-2}, \quad (101)$$

and for $\kappa = \frac{7}{2}$:

$$I(z) = -2(1 + z^2)^{-7/2} \left[\sinh^{-1}(z) - \frac{\pi i}{2} \right] - \frac{16}{15}z(1 + z^2)^{-1} - \frac{4}{3}z(1 + z^2)^{-2} - 2z(1 + z^2)^{-3}. \quad (102)$$

Multiplying equation (99) by the leading coefficient in equation (34), the plasma dispersion function is given by

$$Z_\kappa(\xi) = \frac{1}{(n + \frac{1}{2})^{1/2}} \frac{\Gamma(n + \frac{1}{2})}{\Gamma(n)} \left\{ \sum_{m=1}^n \frac{\Gamma(n + 1 - m)}{\Gamma(n + \frac{3}{2} - m)} z(1 + z^2)^{-m} - \frac{2}{\sqrt{\pi}} (1 + z^2)^{-n-1/2} \left[\sinh^{-1}(z) - \frac{\pi i}{2} \right] \right\}, \quad (103)$$

where $z = \xi/\sqrt{\kappa}$ and $n = 1, 2, 3, \dots$ A more explicit form is obtained by making the substitution given by equation (92). Similar expressions were first obtained by Summers and Thorne¹⁰ using different methods.

10 Solutions for Integer Values: $\kappa = n$

If $\kappa = n$ is a positive integer, a closed form solution can be derived using the identity (61) to obtain a hypergeometric series which terminates; however, this solution lacks mathematical symmetry. It is preferable to proceed

as follows. The hypergeometric series (53) can be written

$$F(1, 2n, n+1, t) = \sum_{m=0}^{\infty} \frac{\Gamma(1+m)}{\Gamma(1)} \frac{\Gamma(2n+m)}{\Gamma(2n)} \frac{\Gamma(n+1)}{\Gamma(n+1+m)} \frac{t^m}{m!} \quad (104)$$

$$= \frac{\Gamma(n+1)}{\Gamma(2n)} \sum_{m=0}^{\infty} \frac{\Gamma(2n+m)}{\Gamma(n+1+m)} t^m \quad (105)$$

$$= \frac{\Gamma(n+1)}{\Gamma(2n)} \sum_{m=n}^{\infty} \frac{\Gamma(m+n)}{\Gamma(m+1)} t^{m-n}. \quad (106)$$

Splitting the sum into two parts

$$F(1, 2n, n+1, t) = \frac{\Gamma(n+1)}{\Gamma(2n)} \left\{ \sum_{m=0}^{\infty} \frac{\Gamma(m+n)}{m!} t^{m-n} - \sum_{m=0}^{n-1} \frac{\Gamma(m+n)}{m!} t^{m-n} \right\} \quad (107)$$

the first part can be summed by the binomial theorem

$$\sum_{m=0}^{\infty} \frac{\Gamma(m+\alpha)}{\Gamma(\alpha)} \frac{z^m}{m!} = (1-z)^{-\alpha}, \quad \alpha \neq 0, -1, -2, \dots, \quad (108)$$

to obtain

$$F(1, 2n, n+1, t) = \frac{\Gamma(n+1)}{\Gamma(2n)} \left\{ \Gamma(n) t^{-n} (1-t)^{-n} - \sum_{m=0}^{n-1} \frac{\Gamma(m+n)}{m!} t^{m-n} \right\}. \quad (109)$$

Substituting $t = (1+iz)/2$ this yields

$$F\left(1, 2n, n+1, \frac{1+iz}{2}\right) = \frac{\Gamma(n+1)\Gamma(n)}{\Gamma(2n)} \left\{ \left(\frac{1+z^2}{4}\right)^{-n} - \sum_{m=0}^{n-1} \frac{\Gamma(m+n)}{\Gamma(n)m!} \left(\frac{1+iz}{2}\right)^{m-n} \right\}. \quad (110)$$

Using the duplication formula (B6) found in Appendix B to evaluate the coefficient on the right hand side, one obtains

$$F\left(1, 2n, n+1, \frac{1+iz}{2}\right) = 2\pi^{1/2} \frac{\Gamma(n+1)}{\Gamma(n+\frac{1}{2})} \left\{ (1+z^2)^{-n} - \frac{1}{4^n} \sum_{m=0}^{n-1} \frac{\Gamma(m+n)}{\Gamma(n)m!} \left(\frac{1+iz}{2}\right)^{m-n} \right\}. \quad (111)$$

Comparing this equation term by term with the symmetry relation [see equation (69)]

$$F\left(1, 2\kappa, \kappa + 1, \frac{1 + iz}{2}\right) + F\left(1, 2\kappa, \kappa + 1, \frac{1 - iz}{2}\right) = 2\sqrt{\pi} \frac{\Gamma(\kappa + 1)}{\Gamma(\kappa + \frac{1}{2})} (1 + z^2)^{-\kappa} \quad (112)$$

it follows that

$$F\left(1, 2n, n + 1, \frac{1 - iz}{2}\right) = 2\pi^{1/2} \frac{\Gamma(n + 1)}{\Gamma(n + \frac{1}{2})} \frac{1}{4^n} \sum_{m=0}^{n-1} \frac{\Gamma(m + n)}{\Gamma(n)m!} \left(\frac{1 + iz}{2}\right)^{m-n} \quad (113)$$

or, equivalently,

$$F\left(1, 2n, n + 1, \frac{1 + iz}{2}\right) = \frac{\pi^{1/2}}{2^{2n-1}} \cdot \frac{n}{\Gamma(n + \frac{1}{2})} \sum_{m=0}^{n-1} \frac{\Gamma(m + n)}{m!} \left(\frac{1 - iz}{2}\right)^{m-n}. \quad (114)$$

Substituting this into equation (56) yields

$$I(z) = \frac{i\pi}{2^{2n-1}} \sum_{m=0}^{n-1} \binom{n + m - 1}{m} \left(\frac{1 - iz}{2}\right)^{m-n}, \quad (115)$$

$n = 1, 2, 3, \dots$ This same result is obtained by using the theory of residues to evaluate the integral (35) along a large semicircle in the lower half plane.² Because (115) is a rational function of z , it immediately gives the analytic continuation of $I(z)$ throughout the complex plane.

Using equation (110) to rewrite (113) in the form

$$F\left(1, 2n, n + 1, \frac{1 + iz}{2}\right) = \frac{\Gamma(n + 1)\Gamma(n)}{\Gamma(2n)} \sum_{m=0}^{n-1} \frac{\Gamma(m + n)}{\Gamma(n)m!} \left(\frac{1 - iz}{2}\right)^{m-n}, \quad (116)$$

it follows from equation (57) that the plasma dispersion function is given by

$$Z_n(\xi) = i \frac{(n - \frac{1}{2})}{n^{1/2}} \frac{\Gamma(n)}{\Gamma(2n)} \sum_{m=0}^{n-1} \frac{\Gamma(m + n)}{m!} \left(\frac{1 - iz}{2}\right)^{m-n}, \quad (117)$$

where $z = \xi/\sqrt{n}$ and $n = 1, 2, 3, \dots$

11 Equivalent Integral Representations

Consider the integral defined for $\text{Im}(z) > 0$ by

$$I(z) = \int_{-\infty}^{\infty} \frac{1}{(1+x^2)^\kappa} \cdot \frac{1}{(x-z)} dx \quad (118)$$

and by analytic continuation for $\text{Im}(z) \leq 0$. By means of some elementary changes of variables, one may derive the following equivalent representations:

$$I(z) = \int_{-\infty}^{\infty} \frac{1}{(1+x^2)^\kappa} \cdot \frac{z}{(x^2-z^2)} dx, \quad (119)$$

$$I(z) = \int_{-\infty}^{\infty} \frac{1}{(\cosh x)^{2\kappa-1}} \cdot \frac{1}{[\sinh(x)-z]} dx, \quad (120)$$

$$I(z) = \int_{-\infty}^{\infty} \frac{1}{(\cosh x)^{2\kappa-1}} \cdot \frac{z}{[\sinh^2(x)-z^2]} dx, \quad (121)$$

$$I(z) = \int_{-\infty}^{\infty} \left(\frac{x^2}{1+x^2} \right)^\kappa \frac{z}{1-z^2x^2} dx, \quad (122)$$

$$I(z) = \int_{-\infty}^{\infty} (\tanh x)^{2\kappa} \frac{z \cosh x}{1-z^2 \sinh^2 x} dx, \quad (123)$$

$$I(z) = z \int_0^\infty t^{\kappa-1/2} (1+t)^{-\kappa} (1-z^2t)^{-1} dt, \quad (124)$$

Thus, by equation (55), all of these integrals are equal to

$$I(z) = i\sqrt{\pi} \frac{\Gamma(\kappa+1/2)}{\Gamma(\kappa+1)} F\left(1, 2\kappa, \kappa+1, \frac{1+iz}{2}\right). \quad (125)$$

In addition, $I(z)$ can be represented as a Laplace transform. If $z = is$ and $\text{Re}(s) > 0$, then one has the identity

$$\frac{1}{x-z} = \frac{i}{s+ix} = i \int_0^\infty e^{-(s+ix)t} dt. \quad (126)$$

Substituting this into (118) and changing the order of integration yields

$$I(is) = i \int_0^\infty f(t) e^{-st} dt, \quad (127)$$

where

$$f(t) = 2 \int_0^\infty \frac{\cos(xt)}{(1+x^2)^\kappa} dx. \quad (128)$$

Using the well known integral representation of the modified Bessel function of the second kind:

$$K_\nu(t) = \frac{\Gamma(\nu + 1/2)2^\nu}{\pi^{1/2}t^\nu} \int_0^\infty \frac{\cos(xt)}{(1+x^2)^{\nu+1/2}} dx, \quad t > 0, \quad (129)$$

it follows that

$$f(t) = \frac{2\sqrt{\pi}}{\Gamma(\kappa)} \left(\frac{t}{2}\right)^{\kappa-1/2} K_{\kappa-1/2}(t), \quad t > 0. \quad (130)$$

This establishes the identity

$$I(z = is) = i\mathcal{L}[f(t)], \quad \text{Re}(s) > 0, \quad (131)$$

where $f(t)$ is defined by the previous equation and \mathcal{L} is the Laplace transform operator.

12 Conclusions

Starting with the definition of the plasma dispersion function in terms of the integral (35), a second order differential equation is derived that immediately allows the dispersion function to be expressed in terms of Gauss' hypergeometric function. Using the well known properties of the hypergeometric function, the many properties of the plasma dispersion function are then derived in a unified manner. A summary of the results is provided in Section 2.

Appendix A: Evaluation of the Integral I_0

By definition

$$I_0 = I(z = i) = \int_{-\infty}^{\infty} \frac{1}{(1+x^2)^\kappa} \cdot \frac{1}{(x-i)} dx. \quad (A1)$$

Multiplying the integrand by $(x+i)/(x+i)$, this becomes

$$I_0 = 2i \int_0^\infty \frac{1}{(1+x^2)^{\kappa+1}} dx, \quad (A2)$$

and with the change of variable $t = x^2$ one obtains

$$I_0 = i \int_0^\infty \frac{t^{-1/2}}{(1+t)^{\kappa+1}} dx. \quad (\text{A3})$$

The remaining integral is the Beta function

$$B(p, q) = \int_0^\infty \frac{t^{p-1}}{(1+t)^{p+q}} dt = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}, \quad (\text{A4})$$

with $p = 1/2$ and $q = \kappa + 1/2$. Therefore, it follows that

$$I_0 = i\sqrt{\pi} \frac{\Gamma(\kappa + 1/2)}{\Gamma(\kappa + 1)}. \quad (\text{A5})$$

The calculation of $I'(z = i)$ is similar.

Appendix B: Alternative Derivation of the Expansion for $|z| > 1$

The purpose of this appendix is to give an independent derivation of the expansion (83) for $|z| > 1$. The starting point is the representation of that function as a Laplace transform. Using the definition

$$K_\nu(x) = \frac{\pi}{2} \frac{I_{-\nu}(x) - I_\nu(x)}{\sin(\pi\nu)}, \quad (\text{B1})$$

together with the series expansion for the modified Bessel function $I_\nu(x)$, one obtains

$$\left(\frac{x}{2}\right)^\nu K_\nu(x) = \frac{\pi}{2\sin(\pi\nu)} \left\{ \sum_{n=0}^{\infty} \frac{1}{\Gamma(n - \nu + 1)n!} \left(\frac{x}{2}\right)^{2n} - \sum_{n=0}^{\infty} \frac{1}{\Gamma(n + \nu + 1)n!} \left(\frac{x}{2}\right)^{2n+2\nu} \right\}, \quad (\text{B2})$$

where $x > 0$ and $\nu > 0$ is not an integer. By a well known theorem,¹¹ the Laplace transform of this function has the asymptotic expansion

$$\mathcal{L}\left\{\left(\frac{t}{2}\right)^\nu K_\nu(t)\right\} \sim \frac{\pi}{2\sin(\pi\nu)} \left\{ \sum_{n=0}^{\infty} \frac{\Gamma(2n+1)}{\Gamma(n-\nu+1)} \frac{1}{n!2^{2n}} \frac{1}{s^{2n+1}} - \sum_{n=0}^{\infty} \frac{\Gamma(2n+2\nu+1)}{\Gamma(n+\nu+1)} \frac{1}{n!2^{2n+2\nu}} \frac{1}{s^{2n+2\nu+1}} \right\}, \quad (\text{B3})$$

where $s \rightarrow \infty$ along any ray $e^{i\theta}$ with $|\theta| < \pi/2$. In fact, the following analysis shows that this series converges in the right half-plane whenever $|s| > 1$ and, therefore, equality holds throughout this region. However, this fact is not needed now. Setting $\nu = \kappa - 1/2$, then from the relations (131) and (56) one obtains

$$\begin{aligned} F\left(1, 2\kappa, \kappa + 1, \frac{1-s}{2}\right) &= \frac{1}{i\sqrt{\pi}} \frac{\Gamma(\kappa+1)}{\Gamma(\kappa+\frac{1}{2})} I(is) \\ &\sim \frac{\pi}{2\sin[(\kappa-\frac{1}{2})\pi]} \frac{\kappa}{\Gamma(\kappa+\frac{1}{2})} \left\{ \sum_{n=0}^{\infty} \frac{\Gamma(2n+1)}{\Gamma(n-\kappa+\frac{3}{2})} \frac{1}{n!2^{2n}} \frac{1}{s^{2n+1}} - \sum_{n=0}^{\infty} \frac{\Gamma(2n+2\kappa)}{\Gamma(n+\kappa+\frac{1}{2})} \frac{1}{n!2^{2n+2\kappa-1}} \frac{1}{s^{2n+2\kappa}} \right\}, \quad (\text{B4}) \end{aligned}$$

where $s \rightarrow \infty$ along any ray $e^{i\theta}$ with $|\theta| < \pi/2$. It will now be shown that this agrees with the expansion (83), that is:

$$\begin{aligned} F\left(1, 2\kappa, \kappa + 1, \frac{1-s}{2}\right) &= \\ &\sqrt{\pi} \frac{\Gamma(\kappa+1)}{\Gamma(\kappa+\frac{1}{2})} [1 + i \tan(\pi\kappa) \text{sgn}[\text{Re}(is)]] (-s^2)^{-\kappa} \left(1 - \frac{1}{s^2}\right)^{-\kappa} \\ &+ \frac{\kappa}{(\kappa-\frac{1}{2})s} \left\{ 1 + \frac{\frac{1}{2}}{\frac{3}{2}-\kappa} \left(\frac{1}{s^2}\right) + \frac{\frac{1}{2} \cdot \frac{3}{2}}{(\frac{3}{2}-\kappa)(\frac{5}{2}-\kappa)} \left(\frac{1}{s^2}\right)^2 + \dots \right\}. \quad (\text{B5}) \end{aligned}$$

The even and odd power terms are compared separately. First the even terms. Using the duplication formula

$$\Gamma(2z) = \pi^{-1/2} 2^{2z-1} \Gamma(z) \Gamma\left(z + \frac{1}{2}\right) \quad (\text{B6})$$

and the binomial theorem, the even terms in (B4) can be summed as follows

$$\begin{aligned}
 & - \frac{\pi}{\sin[(\kappa - \frac{1}{2})\pi]} \frac{\kappa}{\Gamma(\kappa + \frac{1}{2})} \sum_{n=0}^{\infty} \frac{1}{n!} \frac{\Gamma(2n + 2\kappa)}{\Gamma(n + \kappa + \frac{1}{2})} \frac{1}{2^{2n+2\kappa-1} s^{2n+2\kappa}} \\
 & = \frac{\sqrt{\pi}}{\cos(\kappa\pi)} \frac{\kappa\Gamma(\kappa)}{\Gamma(\kappa + \frac{1}{2})} \sum_{n=0}^{\infty} \frac{\Gamma(n + \kappa)}{\Gamma(\kappa)n!} \frac{1}{s^{2n+2\kappa}} \\
 & = \frac{\sqrt{\pi}}{\cos(\kappa\pi)} \frac{\kappa\Gamma(\kappa)}{\Gamma(\kappa + \frac{1}{2})} s^{-2\kappa} \left(1 - \frac{1}{s^2}\right)^{-\kappa}. \quad (\text{B7})
 \end{aligned}$$

This will now be compared with the even terms in (B5):

$$\sqrt{\pi} \frac{\Gamma(\kappa + 1)}{\Gamma(\kappa + \frac{1}{2})} [1 + i \tan(\pi\kappa) \text{sgn}[\text{Re}(is)]] (-s^2)^{-\kappa} \left(1 - \frac{1}{s^2}\right)^{-\kappa}. \quad (\text{B8})$$

It is easy to show that

$$\frac{(-s^2)^{-\kappa}}{s^{-2\kappa}} = \begin{cases} e^{+i\pi\kappa} & \text{if } 0 < \arg(s) < \pi/2 \\ e^{-i\pi\kappa} & \text{if } -\pi/2 < \arg(s) \leq 0, \end{cases} \quad (\text{B9})$$

therefore,

$$\frac{s^{-2\kappa}}{\cos(\kappa\pi)} = [1 + i \tan(\pi\kappa) \text{sgn}[\text{Re}(is)]] (-s^2)^{-\kappa}, \quad (\text{B10})$$

which establishes the equality of (B7) and (B8), that is, for the even power terms in (B4) and (B5). Similarly, using the duplication formula (B6) the odd terms in (B4) are easily shown to be equal to the odd terms in (B5). This gives an independent proof of (83) in the upper half plane which, by means of the symmetry relations, extends to the whole plane.

References

1. D. Summers and R. M. Thorne, “The modified plasma dispersion function,” *Phys. Fluids B* 3, 1835-1847, 1991.
2. Y. F. Chateau and N. Meyer-Vernet, “Electrostatic Noise in Non-Maxwellian Plasmas: Generic Properties and Kappa Distributions,” *J. Geophys. Res.* 96A, 5825-5836, 1991.

3. R. L. Mace and M. A. Hellberg, “A dispersion function for plasmas containing superthermal particles,” *Phys. Plasmas* 2, 2098-2109, 1995.
4. N. A. Krall and A. W. Trivelpiece, *Principles of Plasma Physics*, San Francisco Press, 1973, Chapter 8.
5. D. G. Swanson, *Plasma Waves*, Academic Press, 1989, page 129.
6. B. D. Fried and S. D. Conte, *The Plasma Dispersion Function*, Academic Press, 1961.
7. M. Abramowitz and I. A. Stegun, eds., *Handbook of Mathematical Functions*, 10th printing, 1972, Chapter 15.
8. H. Bateman and A. Erdélyi, *Higher Transcendental Functions*, McGraw-Hill, 1953, Vol I, page 65, equation 28.
9. G. E. Andrews, R. Askey, and R. Joy, *Special Functions*, Cambridge University Press, 1999, page 128, equation 3.1.12.
10. D. Summers, R. M. Thorne and H. Matsumoto, “Evaluation of the modified plasma dispersion function for half-integral indices,” *Phys. Plasmas* 3, 2496-2501, 1996.
11. G. Doetsch, *Introduction to the Theory and Application of the Laplace Transformation*, Springer-Verlag, 1974, page 228.

Author’s Note

This work was performed in the year 2000 during my tenure as a Ph.D. student at the Courant Institute of Mathematical Sciences, New York University.

REPORT DOCUMENTATION PAGE				<i>Form Approved</i> <i>OMB No. 0704-0188</i>	
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>					
1. REPORT DATE (DD-MM-YYYY)		2. REPORT TYPE		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSORING/MONITOR'S ACRONYM(S)	
				11. SPONSORING/MONITORING REPORT NUMBER	
12. DISTRIBUTION/AVAILABILITY STATEMENT					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19b. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (Include area code)